

Corrections

Stoichiometry of the Topa Quinone Biogenesis Reaction in Copper Amine Oxidases, by Christy E. Ruggiero and David M. Dooley*, Volume 38, Number 10, March 9, 1999, pages 2892–2898.

Page 2893. The experimental conditions for the measurement of the production of H₂O₂ were inaccurately reported. Thus, in column 2, lines 13–22 should be replaced with the following: This molar extinction coefficient was determined by H₂O₂ titration at the exact pH and concentration conditions used in the peroxide production experiments (final dye solution = 20 units/mL HRP, 2 mM vanillic acid, 1 mM 4-aminoantipyrine in chelexed Hepes buffer, pH 6.8). The TPQ biogenesis reaction was initiated by the addition of copper (~4-fold molar excess over subunit) to a cuvette containing the dye solution and the apo, unprocessed enzyme (final concentration approximately 15 μM) in a reaction volume of 0.8 mL.

BI9950802

10.1021/bi9950802

Published on Web 07/02/1999

Backbone Dynamics of Apocytochrome *b*₅ in Its Native, Partially Folded State, by Shibani Bhattacharya, Christopher J. Falzone, and Juliette T. J. Lecomte*, Volume 38, Number 8, February 23, 1999, pages 2577–2589.

Page 2588. In Supporting Information, the vertical axis of Figure S1 is incorrect. The corrected figure is available free of charge via the Internet at <http://pubs.acs.org>.

BI995081U

10.1021/bi995081u

Published on Web 07/02/1999

Primary Charge Separation Routes in the BChl:BPhe Heterodimer Reaction Centers of *Rhodobacter sphaeroides*, by Marion E. van Brederode, Ivo H. M. van Stokkum, Evaldas Katilius, Frank van Mourik, Michael R. Jones, and Rienk van Grondelle*, Volume 38, Number 23, June 8, 1999, pages 7545–7555.

Page 7554. The following paragraph was omitted.

SUPPORTING INFORMATION AVAILABLE

Typical datasets of transient absorbance difference spectra of the HM202L RC recorded at 77 K. This material is available free of charge via the Internet at <http://pubs.acs.org>.

BI9950793

10.1021/bi9950793

Published on Web 07/02/1999