

# Hydrogen Bond Switching among Flavin and Amino Acids Determines the Nature of Proton-Coupled Electron Transfer in BLUF Photoreceptors

Tilo Mathes,<sup>\*,†,‡</sup> Jingyi Zhu,<sup>†</sup> Ivo H. M. van Stokkum,<sup>†</sup> M. L. Groot,<sup>†</sup> Peter Hegemann,<sup>‡</sup> and John T. M. Kennis<sup>\*,†</sup>

<sup>†</sup>Biophysics Group, Department of Physics and Astronomy, Faculty of Sciences, Vrije Universiteit, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands

<sup>‡</sup>Institut für Biologie, Experimentelle Biophysik, Humboldt Universität zu Berlin, Invalidenstraße 42, D-10115 Berlin, Germany

**Supporting Information** 

**ABSTRACT:** BLUF domains are flavin-binding photoreceptors that can be reversibly switched from a dark-adapted state to a light-adapted state. Proton-coupled electron transfer (PCET) from a conserved tyrosine to the flavin that results in a neutral flavin semiquinone/tyrosyl radical pair constitutes the photoactivation mechanism of BLUF domains. Whereas in the dark-adapted state PCET occurs in a sequential fashion where electron transfer precedes proton transfer, in the light-adapted state the same radical pair is formed by a concerted mechanism. We propose that the altered nature of the PCET process results from a hydrogen bond switch between the flavin and its surrounding amino acids that preconfigures the system for proton transfer. Hence, BLUF domains represent an attractive biological model system to investigate and understand PCET in great detail.



#### **SECTION:** Biophysical Chemistry

**B** iological photoreceptors consist of a pigment to absorb light of a certain wavelength region bound to a protein that reacts upon photoinduced chemical or structural changes of the pigment to subsequently modulate its interaction with a signaling partner. Most photoreceptors reversibly switch from a dark state to a light-adapted or signaling state upon light absorption. The light-adapted state, a molecular ground state that activates downstream effectors and hence causes a biological reaction, is metastable and recovers thermally to the dark-adapted state.

BLUF photoreceptors are small flavin containing proteins<sup>1</sup> involved in phototaxis,<sup>2</sup> photosynthetic gene regulation,<sup>3</sup> and virulence<sup>4</sup> and have also been found coupled to various enzyme domains.<sup>2,5–9</sup> The latter have gained immense attention as so-called optogenetic tools.<sup>9–13</sup> The essential components of the photosensory apparatus consist of the isoalloxazine moiety of the flavin, a conserved tyrosine and glutamine side-chain. Figure 1 shows the X-ray structure of the Slr1694 BLUF domain from *Synechocystis sp.* PCC 6803<sup>14</sup> (also known as SyPixD) with the flavin binding pocket highlighted. In the BLUF dark state, after excitation with blue light an electron is transferred within a few tens of picoseconds from the tyrosine to the flavin, resulting in a spectroscopically observable FAD<sup>6–</sup> anion semiquinone radical.<sup>15,16</sup> Subsequently FAD<sup>6–</sup> is protonated, most likely by the same tyrosine, to form a neutral flavin semiquinone (FADH<sup>6</sup>)/tyrosyl (TyrO<sup>6</sup>) radical pair. Thus, proton-coupled electron transfer (PCET) underlies



**Figure 1.** Dark- and light-adapted states of Slr1694. A and B show the presumed respective tyrosine/glutamine/flavin hydrogen bond network. The absorbance of both states is presented in C.

BLUF photoactivation, whereby electron transfer precedes proton transfer by picoseconds. During radical recombination a rearrangement of the hydrogen bond network around the flavin takes place leading to a 10-15 nm red shift in the visible absorbance of the flavin as shown in Figure 1C,<sup>3</sup> as well as to a downshift of the carbonyl vibrations of the flavin.<sup>17</sup> The resulting hydrogen bond-switched state is formed within less than 100 ps and remains stable for about 10 decades of time and is therefore considered as the light-adapted or signaling state of the BLUF domain.<sup>15</sup> The details of this hydrogen bond switch are still under considerable discussion since crystal and

Received:December 1, 2011Accepted:December 26, 2011Published:December 26, 2011

NMR structures gave contradicting findings about dark and light-adapted states with respect to glutamine orientation.<sup>6,14,18–24</sup> A likely mechanism involves rotation of the glutamine amide.<sup>15,18</sup> However, recent theoretical calculations introduced new models for the hydrogen bond switch, involving tautomerization with or without rotation of the glutamine side-chain.<sup>25,26</sup>

Because of their unique primary photochemistry and their small size, BLUF domains represent an ideal naturally occurring model system to investigate PCET in great molecular detail. PCET is a key step in solar energy conversion that still requires mechanistic elucidation, especially in photosystem II where a key tyrosine mediates redox processes between P680 and the oxygen-evolving complex.<sup>27–29</sup> With BLUF domains we have an easily accessible protein-chromophore complex whose photoreactivity is determined by well-defined structural constraints and controlled redox partner interactions through hydrogen bonding. Deeper understanding of these evolutionary optimized environments is of major importance to design efficient artificial systems for solar energy conversion.<sup>30</sup>

In this study we investigated the photochemistry of the lightadapted ground state of Slr1694 from Synechocystis sp. PCC 6803. We chose this protein because it is available in milligram amounts of high purity and its light-driven PCET processes are kinetically well-resolved in the dark-adapted state to signalingstate forward reaction.<sup>15,16</sup> To keep Slr1694 in the light-adapted state conveniently, we used the slow-photocycling W91F mutant of Slr1694, which exhibits a 50-fold slower recovery to the dark state as compared to wild type (223 s vs 4 s, for W91F and wild type, respectively).<sup>31</sup> Background illumination with a blue LED then induces formation of the 12 nm redshifted light-adapted state at near 100% population. It is important to note that the ultrafast light-induced dynamics of the Slr1694-W91F dark state remain essentially identical to those of the wild type.<sup>31</sup> With the information on the ultrafast electronic and vibrational dynamics obtained on this system here we show that in the light-adapted state the same neutral flavin semiguinone tyrosyl radical pair is formed as in the darkadapted state reaction. Most importantly we show that its formation by proton coupled electron transfer is transformed from a sequential fashion in the dark-adapted state into a concerted fashion in the light-adapted state, hence making it an ideal model system to study PCET. These changes in photodynamics furthermore strongly support one of the highly debated molecular models of both states as postulated earlier<sup>16</sup> and will be discussed below.

Ultrafast visible absorption dynamics were recorded by transient pump/probe absorption spectroscopy on light-adapted Slr1694-W91F in  $H_2O$  and  $D_2O$  buffer with excitation at 400 nm on an experimental setup described previously.<sup>32</sup> Data treatment and analysis is described in detail in the Supporting Information. The obtained data was sufficiently described by global analysis<sup>33</sup> using a sequential model with four lifetimes.

The evolutionary-associated difference spectra (EADS) are displayed in Figure S2, whereas kinetics at selected wavelengths are shown in Figure 2A. Formation of a flavin semiquinone intermediate is clearly observed by the rising absorption between 550 and 650 nm. The signal is overlaid with the excited state absorption (ESA) at 520 nm and above 630 nm and partly diminished by stimulated emission (SE). This is best demonstrated by the change in absorption at 556 nm (Figure 2A). The initial negative absorption by SE is quickly recovered



Letter

**Figure 2.** Time traces of the light-adapted state absorption changes after 400 nm excitation at selected wavelengths together with their fit. The traces at 556 nm (A) represent SE of the flavin, which is compensated within the first 2 ps by semiquinone absorption. Characteristic absorption of the flavin neutral semiquinone is observed at 617 nm, which is biased by overlap with SE within the first 10 ps. This leads to an apparent slowed down formation of the semiquinone in D<sub>2</sub>O due to a longer lived excited state lifetime in D<sub>2</sub>O (Figure 3A). At delays greater than 10 ps the absorbance change is significantly slowed down in D<sub>2</sub>O at both wavelengths. The IR traces (B) recorded in D<sub>2</sub>O buffer at 1691 and 1628 cm<sup>-1</sup> display the decay of the carbonyl stretch of the flavin and a tyrosine species, respectively. The time axis is linear until 1 (A) and 10 ps (B) and logarithmic after indicated by the skewed lines.

in an H/D isotope independent manner. After reaching a maximum, the signal, now fully belonging to the FAD neutral semiquinone, decays again. This decay is slowed down in the deuterated species. The corresponding EADS (Figure S2A/B) represent mixtures of FAD excited state, FAD neutral semiquinone, and a nondecaying rather featureless spectrum with slightly rising absorbance toward the red edge of the spectral window and a bleach in the flavin ground state absorption region. This spectrum contributes very little to the overall spectral evolution and is therefore difficult to interpret. It might be attributed to solvated electron generation by two-photon ionization<sup>34</sup> observed previously in PYP<sup>35</sup> or a minor fraction of triplet species.<sup>36</sup>

To obtain the pure spectra from this mixture we performed target analysis<sup>33</sup> using a simple compartmental model (Figure 3A). A parallel reaction (green) is included to attribute for the putative photoionization product, which is instantly formed and remains constant on the time scale of the experiment. The mixture of excited state and semiguinone spectra is addressed by introducing a minor heterogeneity of the excited state. Only one of the two spectrally identical excited state species forms the neutral semiquinone. The resulting species-associated difference spectra (SADS) are displayed in Figure 3B. The excited state spectra are as a result of the imposed constraint identical and decay with 1 and 5 ps in H<sub>2</sub>O. The FAD excitedstate lifetime in dark-adapted Slr1694 has multiphasic lifetimes of 7, 40, and 180 ps.<sup>15</sup> Thus, the FAD excited-state lifetime is significantly shortened with respect to that observed in the dark-adapted state,<sup>15,16</sup> as previously observed for the lightadapted states of the Tll0078 and AppA BLUF domains.<sup>37,38</sup> The spectra are highly similar to the EADS and show the ground-state bleach (GSB) of the light-adapted state at 452 nm as well as ESA at 510 nm and from 600 to 725 nm. Stimulated emission (SE) is observed at 560 nm. The fast disappearance corresponds to the formation of the FADH<sup>•</sup> neutral semi-



Figure 3. Target analysis of vis and IR absorption data using the model in A with the corresponding lifetimes in  $H_2O$  ( $D_2O$ ). B and D show the SADS of  $H_2O$  and  $D_2O$  visible absorption data sets. The black and red spectra ideally represent flavin excited state and neutral flavin semiquinone, respectively. The green, nondecaying spectrum contributes little to the data sets and is tentatively assigned to a two-photon product or a minor amount of flavin triplet species. C shows the SADS of the IR data set in  $D_2O$ , obtained by simultaneous analysis of both vis and IR data. The black spectrum is identified as a flavin excited state whereas the red spectrum features a new bleach at around 1628 cm<sup>-1</sup>, which is attributed to the tyrosine radical pair partner.

quinone (red), which lives for 66 ps. The species is formed with an efficiency of about 40% as judged by the loss of GSB. The FADH<sup>•</sup> spectrum shows the same GSB and additional absorptive features between 550 and 625 nm, representing the neutral radical absorption.<sup>15</sup> The nondecaying spectrum (green) is identical to the spectrum from the sequential analysis. In D<sub>2</sub>O we observe similar spectra with different lifetimes. The excited states decay with 1.1 and 6.9 ps, which represents a KIE of 1.1 and 1.4, respectively. Thus, we observe a significant kinetic isotope effect for the slow fraction of FAD\* decay (5 vs 6.9 ps). The red spectrum is formed with an efficiency of about 50% and represents the absorption of a neutral flavin semiquinone FADH<sup>•</sup>. The spectrum decays with 140 ps and shows a KIE of about 2.1.

Strikingly, the spectra and the lifetimes of the FADH<sup>•</sup> neutral semiquinone are practically identical to the dark state intermediate of this particular Slr1694 mutant, in H<sub>2</sub>O as well as  $D_2O$ .<sup>31</sup> Therefore the process of radical recombination, which leads to the red-shifted, oxidized BLUF state, must be highly similar to that of the dark state reaction. This observation strongly suggests that, in the light adapted state of BLUF domains, the same FADH<sup>•</sup> intermediate with regard to hydrogen-bond pattern and glutamine orientation is formed as in the dark state reaction.

To confirm the identity of the dark- and light-adapted state intermediates as suggested above, we identified the corresponding radical pair partner of the neutral flavin semiquinone by femtosecond infrared spectroscopy. To prevent the large underlying absorption of the OH bending mode of water we measured the spectral dynamics between 1460 and 1740 cm<sup>-1</sup> in D<sub>2</sub>O. The most prominent signals are observed at 1691 cm<sup>-1</sup> (Figure 2B) and 1549 cm<sup>-1</sup> (Figure S3), which represent the carbonyl stretch and the C==N out-of-phase vibration of the flavin, respectively. Both signals decay completely with the same kinetics on the observed time scale. The induced absorption of the carbonyl stretch at  $1674 \text{ cm}^{-1}$  and the bleach of the xylene ring vibration of the flavin at 1647  $\text{cm}^{-1}$  decay in parallel (Figure S3). The corresponding induced absorption of the latter at 1628  $\text{cm}^{-1}$  (Figure 2B), however, rises initially but then decays quickly within 5 ps and becomes negative, indicating a ground-state bleach of a molecular group other than the flavin. The negative absorption then decays within the observed time scale. A bleach at around 1620 cm<sup>-1</sup> was previously assigned to a tyrosine ring vibration and interpreted as a loss of an electron and a proton in formation of a tyrosyl radical.<sup>16</sup> The corresponding EADS of the IR measurements (Figure S2C) are highly similar to the SADS (Figure 3C). Both show a prominent carbonyl bleach of the flavin cofactor at around 1690  $\text{cm}^{-1}$  and the C=N vibration at 1550  $\text{cm}^{-1}$  as previously observed.<sup>16</sup> The former is downshifted relative to the dark state by  $\sim 10 \text{ cm}^{-1}$  and indicates stronger hydrogen bonding as previously observed for the light-adapted state by steady state FTIR.<sup>17</sup> The occurrence of a previously identified tyrosine signature at 1628 cm<sup>-1</sup> clearly demonstrates that the radical pair consists of the flavin and the close-by tyrosine, identical to the dark state intermediate.<sup>16</sup> The IR absorbance difference spectra of FAD\* and FADH reported in Figure 3C may be compared to the corresponding spectra in dark-adapted BLUF domains.<sup>16</sup>

Upon dark-state excitation, PCET in Slr1694 and most likely in BLUF domains in general occurs in a sequential fashion where proton transfer lags electron transfer by picoseconds.<sup>15,16,39</sup> Here, we have shown that a FADH<sup>•</sup> neutral semiguinone similar or identical to that formed in the darkstate reaction, is very rapidly produced by electron and proton transfer in 1 ps without any observable anionic FAD<sup>•-</sup> radical intermediate. Hence, PCET occurs in a rather concerted fashion after light-adapted state excitation in BLUF domains. In general, a concerted mechanism features a lower activation barrier since it does not involve the charged high-energy intermediates that result from separated proton and electron transfer.<sup>40</sup> The reconfiguration of the hydrogen-bond network around the FAD chromophore in the BLUF domain very likely is responsible for the altered nature of the PCET process. To facilitate the obviously more efficient proton tunnelling in the light-adapted state of BLUF, the molecular interactions need to be preconfigured for proton transfer either by shortening the proton transfer distance or by establishing an indirect proton conducting wire. Similar to the AppA and Tll0078 BLUF domains, the heterogeneity in the excited-state decay in Slr1694-W91F is significantly reduced in the light-adapted state.<sup>37,38</sup> This supports the general assumption that the coordination of the flavin becomes tighter, more rigid and less heterogeneous in the light-adapted state. After femtosecond excitation the light-adapted excited state decays rapidly with an H/D isotope independent lifetime of 1 ps to form the neutral flavin radical. Because we do not observe any electron transfer intermediate like an anionic semiquinone here, the proton transfer clearly must occur in less than 1 ps, which can be explained by the increased basicity of the nearby N5 atom in the excited state of flavin.<sup>41</sup> It should be noted that the time scale of this reaction is in the range of skeletal motions ( $\sim$ 300 fs) that finally mediate excited state proton transfer as previously observed in GFP.<sup>42</sup> Hence, the proton transfer rate observed here has been pushed close to the physical limit by the reconfiguration of the hydrogen bond network around the flavin in the light adapted state.

The exact configuration of the hydrogen bond network and orientation of the glutamine side chain in the light-adapted state, which determine the changed reactivity we observe here, are still under heavy debate.<sup>16,18,19,25,26,43-45</sup> Recent NMR and FTIR studies showed that an unusually strong hydrogen bond from the tyrosine hydroxyl group to the glutamine is formed in the BLUF light-adapted state.<sup>45,46</sup> The molecular model of the light-adapted state that we proposed earlier (Figure 4A,



**Figure 4.** Models for PCET in the light state of Slr1694. The reaction involving concerted electron/proton transfer (CEPT) as proposed here is displayed in (A), with CEPT from Y8 to FAD taking place with a time constant of 1 ps after photoexcitation, see the main text for details. Alternative starting geometries (B and C) might involve the tautomeric forms observed in computational models and might involve direct H-atom transfer (C).

identical to the light state configuration in ref 15) provides the basis for such a strong hydrogen bond between tyrosine and glutamine. On the basis of the kinetic information on PCET in the light state of BLUF provided here, we propose that the formal hydrogen atom transfer is occurring by a concerted electron and proton transfer with a time constant of 1 ps, with direct electron transfer from the tyrosine and proton transfer by a Grotthus-like mechanism along the amide group of the glutamine to FAD N5, to result in formation of a FADH<sup>•</sup>-Y8<sup>•</sup> radical pair. Due to the presumably tighter coordination of the flavin in the light-adapted state, the proton is instantly transferred. An intermediary imidic glutamine side chain, which might form upon transfer of the tyrosine hydroxyl proton to the amide carbonyl and release of an amide nitrogen proton to N5, instantly retautomerizes to the energetically more favorable amide form (Figure 4A, right). Note that the resulting molecular configuration between Y8°, Q50, and FADH<sup>•</sup> (Figure 4A, right) matches that proposed for the FADH<sup>•</sup>-Y8<sup>•</sup> radical pair intermediate of the dark-state reaction of the same protein,<sup>15</sup> which is consistent with our observation that these two radical pairs have similar spectra, the same lifetime of 66 ps and the same kinetic isotope effect of 2.1 upon H/D exchange. Yet, the details of the binding pocket with respect to side chain configuration remain to be further characterized and cannot be assigned here completely. A direct hydrogen atom transfer from the tyrosine seems unlikely concerning the already hydrogen bonded N5 of the flavin and the rather large distance of 4-5 Å to the flavin.

In QM/MM studies of Domratcheva et al.<sup>26</sup> and Sadeghian et al.<sup>25,47</sup> and spectroscopic studies by Stelling and co-workers<sup>43</sup> alternative models for the BLUF light-adapted state were proposed that involve tautomeric forms of Q50, as indicated in Figure 4, parts B and C. Although a tautomeric form may energetically be unfavorable for the long-lived signaling state in BLUF<sup>48</sup> we will discuss these models with respect to strong hydrogen bonding as observed by FTIR<sup>46</sup> and fast PCET observed in this study. In Domratcheva's model (Figure 4B), the imidic nitrogen of the glutamine side chain receives a hydrogen bond from Q50 and donates a hydrogen bond to flavin N5.<sup>26</sup> In such a molecular configuration, a concerted mechanism with a very short proton relay involving only the nitrogen atom of the amide group of Q50 would present a very efficient proton transfer path consistent with the experimental results presented here. However, this molecular configuration for the light-adapted state is difficult to reconcile with the presence of an unusually strong hydrogen bond between Y8 and Q50:46 in general stronger H-bonds are formed to the oxygen rather than to the nitrogen of an amide or an imidic acid due to the higher electronegativity of oxygen. Hence, we consider this model less likely.

In Sadeghian's model the imidic acid oxygen of the glutamine side chain receives a hydrogen bond from Q50 and donates a hydrogen bond to flavin N5 (Figure 4C).<sup>25</sup> In this molecular configuration, a direct H-atom transfer from Y8 to FAD may readily occur because the H-bond of the imidic Q50 side chain to N5 is easily removed by tautomerization to the energetically favorable amide form. However, this molecular model is not consistent with formation of an unusually strong hydrogen bond between Y8 and the imidic oxygen of Q50: strong hydrogen bonds acquire significant covalent bond character,<sup>49</sup> which would destabilize the tautomeric form of the glutamine side chain even further, rendering this model for the BLUF light-adapted state less likely.

An interesting result from our experiments is the H/Disotope dependent decay of a fraction of FAD excited-states, in 5 ps in  $H_2O$  and 6.9 ps in  $D_2O$  (Figure 3A). This fraction does not form an obvious proton associated product such as a flavin neutral semiquinone, but rather returns to the FAD ground state directly. The isotope-dependent excited-state decay might be due to a reconfiguration of the H-bond network in the excited state. Most likely FAD N5 attracts an H-bond stronger due to its increased basicity in the singlet excited state.<sup>41</sup> The observation of this process in only a fraction of the proteins shows that a certain degree of ground-state heterogeneity exists in the light-adapted state, with a protein configuration in which the electron transfer is inefficient, probably due to an unfavorable tyrosine/flavin orientation.

In summary, our study demonstrates that an intricate biological hydrogen bond network tunes the photoinduced electron/proton transfer of a flavin/protein complex from a sequential to a concerted reaction. Since the BLUF domain can be easily switched between both reactivities it clearly represents a most powerful model system for further studies on protoncoupled electron transfer in biological systems.

## ASSOCIATED CONTENT

#### Supporting Information

Experimental procedures for sample preparation, spectroscopy and data analysis as well as supplementary data. This material is available free of charge via the Internet at http://pubs.acs.org.

#### AUTHOR INFORMATION

#### **Corresponding Author**

\*E-mail: t.mathes@vu.nl (T.M.); j.t.m.kennis@vu.nl (J.T.M.K.).

#### ACKNOWLEDGMENTS

T.M. was supported by the Federation of European Biochemical Societies (FEBS), the LaserLab Europe access program (LCVU-1618). The authors acknowledge a NWO-CW ECHO grant to J.T.M.K. and the DFG FOR1279 to P.H.

# REFERENCES

(1) Gomelsky, M.; Klug, G. BLUF: a novel FAD-binding domain involved in sensory transduction in microorganisms. *Trends Biochem. Sci.* **2002**, *27*, 497.

(2) Iseki, M.; Matsunaga, S.; Murakami, A.; Ohno, K.; Shiga, K.; Yoshida, K.; Sugai, M.; Takahashi, T.; Hori, T.; Watanabe, M. A bluelight-activated adenylyl cyclase mediates photoavoidance in Euglena gracilis. *Nature* **2002**, *415*, 1047.

(3) Masuda, S.; Bauer, C. E. AppA is a blue light photoreceptor that antirepresses photosynthesis gene expression in Rhodobacter sphaeroides. *Cell* **2002**, *110*, 613.

(4) Mussi, M. A.; Gaddy, J. A.; Cabruja, M.; Arivett, B. A.; Viale, A. M.; Rasia, R.; Actis, L. A. The opportunistic human pathogen Acinetobacter baumannii senses and responds to light. *J. Bacteriol.* **2010**, *192*, 6336.

(5) Rajagopal, S.; Key, J. M.; Purcell, E. B.; Boerema, D. J.; Moffat, K. Purification and initial characterization of a putative blue light-regulated phosphodiesterase from Escherichia coli. *Photochem. Photobiol.* **2004**, *80*, 542.

(6) Barends, T. R. M.; Hartmann, E.; Griese, J. J.; Beitlich, T.; Kirienko, N. V.; Ryjenkov, D. A.; Reinstein, J.; Shoeman, R. L.; Gomelsky, M.; Schlichting, I. Structure and mechanism of a bacterial light-regulated cyclic nucleotide phosphodiesterase. *Nature* **2009**, *459*, 1015.

(7) Ryu, M. H.; Moskvin, O. V.; Siltberg-Liberles, J.; Gomelsky, M. Natural and engineered photoactivated nucleotidyl cyclases for optogenetic applications. *J. Biol. Chem.* **2010**, *285*, 41501.

(8) Tschowri, N.; Busse, S.; Hengge, R. The BLUF-EAL protein YcgF acts as a direct anti-repressor in a blue-light response of Escherichia coli. *Genes Dev.* **2009**, *23*, 522.

(9) Stierl, M.; Stumpf, P.; Udwari, D.; Gueta, R.; Hagedorn, R.; Losi, A.; Gärtner, W.; Petereit, L.; Efetova, M.; Schwarzel, M.; et al. Light modulation of cellular cAMP by a small bacterial photoactivated adenylyl cyclase, bPAC, of the soil bacterium Beggiatoa. *J. Biol. Chem.* **2011**, 286, 1181.

(10) Bucher, D.; Buchner, E. Stimulating PACalpha increases miniature excitatory junction potential frequency at the Drosophila neuromuscular junction. *J. Neurogenet.* **2009**, *23*, 220.

(11) Nagahama, T.; Suzuki, T.; Yoshikawa, S.; Iseki, M. Functional transplant of photoactivated adenylyl cyclase (PAC) into Aplysia sensory neurons. *Neurosci. Res.* **2007**, *59*, 81.

(12) Weissenberger, S.; Schultheis, C.; Liewald, J. F.; Erbguth, K.; Nagel, G.; Gottschalk, A. PACalpha - an optogenetic tool for in vivo manipulation of cellular cAMP levels, neurotransmitter release, and behavior in Caenorhabditis elegans. *J. Neurochem.* **2011**, *116*, 616.

(13) Schröder-Lang, S.; Schwärzel, M.; Seifert, R.; Strünker, T.; Kateriya, S.; Looser, J.; Watanabe, M.; Kaupp, U. B.; Hegemann, P.; Nagel, G. Fast manipulation of cellular cAMP level by light in vivo. *Nat. Methods* **2007**, *4*, 39.

(14) Yuan, H.; Anderson, S.; Masuda, S.; Dragnea, V.; Moffat, K.; Bauer, C. Crystal structures of the Synechocystis photoreceptor Slr1694 reveal distinct structural states related to signaling. *Biochemistry* **2006**, *45*, 12687.

(15) Gauden, M.; Stokkum, I. H. M. v.; Key, J. M.; Lührs, D. C.; Grondelle, R. v.; Hegemann, P.; Kennis, J. T. M. Hydrogen-bond switching through a radical pair mechanism in a flavin-binding photoreceptor. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 10895.

(16) Bonetti, C.; Mathes, T.; van Stokkum, I. H. M.; Mullen, K. M.; Groot, M.-L.; van Grondelle, R.; Hegemann, P.; Kennis, J. T. M. Hydrogen Bond Switching among Flavin and Amino Acid Side Chains in the BLUF Photoreceptor Observed by Ultrafast Infrared Spectroscopy. *Biophys. J.* **2008**, *95*, 4790.

(17) Masuda, S.; Hasegawa, K.; Ishii, A.; Ono, T.-a. Light-induced structural changes in a putative blue-light receptor with a novel FAD binding fold sensor of blue-light using FAD (BLUF); Slr1694 of synechocystis sp. PCC6803. *Biochemistry* **2004**, *43*, 5304.

(18) Anderson, S.; Dragnea, V.; Masuda, S.; Ybe, J.; Moffat, K.; Bauer, C. Structure of a novel photoreceptor, the BLUF domain of AppA from Rhodobacter sphaeroides. *Biochemistry* **2005**, *44*, 7998.

(19) Jung, A.; Reinstein, J.; Domratcheva, T.; Shoeman, R.; Schlichting, I. Crystal Structures of the AppA BLUF Domain Photoreceptor Provide Insights into Blue Light-mediated Signal Transduction. J. Mol. Biol. 2006, 362, 717.

(20) Jung, A.; Domratcheva, T.; Tarutina, M.; Wu, Q.; Ko, W.-H.; Shoeman, R. L.; Gomelsky, M.; Gardner, K. H.; Schlichting, I. Structure of a bacterial BLUF photoreceptor: insights into blue lightmediated signal transduction. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 12350.

(21) Kita, A.; Okajima, K.; Morimoto, Y.; Ikeuchi, M.; Miki, K. Structure of a cyanobacterial BLUF protein, Tll0078, containing a novel FAD-binding blue light sensor domain. *J. Mol. Biol.* **2005**, *349*, 1. (22) Wu, Q.; Gardner, K. H. Structure and Insight into Blue Light-

Induced Changes in the BlrP1 BLUF Domain. *Biochemistry* 2009, 48, 2620.

(23) Wu, Q.; Ko, W.-H.; Gardner, K. H. Structural requirements for key residues and auxiliary portions of a BLUF domain. *Biochemistry* **2008**, *47*, 10271.

(24) Grinstead, J. S.; Hsu, S. T.; Laan, W.; Bonvin, A. M.; Hellingwerf, K. J.; Boelens, R.; Kaptein, R. The solution structure of the AppA BLUF domain: insight into the mechanism of light-induced signaling. *ChemBioChem* **2006**, *7*, 187.

(25) Sadeghian, K.; Bocola, M.; Schütz, M. A conclusive mechanism of the photoinduced reaction cascade in blue light using flavin photoreceptors. J. Am. Chem. Soc. 2008, 130, 12501.

(26) Domratcheva, T.; Grigorenko, B. L.; Schlichting, I.; Nemukhin, A. V. Molecular models predict light-induced glutamine tautomerization in BLUF photoreceptors. *Biophys. J.* **2008**, *94*, 3872.

(27) Barry, B. A. Proton coupled electron transfer and redox active tyrosines in Photosystem II. J. Photochem. Photobiol. B: Biol. **2011**, 104, 60–71.

(28) Huynh, M. H. V.; Meyer, T. J. Proton-coupled electron transfer. *Chem. Rev.* **2007**, *107*, 5004.

(29) Moore, G. F.; Hambourger, M.; Kodis, G.; Michl, W.; Gust, D.; Moore, T. A.; Moore, A. L. Effects of protonation state on a tyrosinehistidine bioinspired redox mediator. *J. Phys. Chem. B* **2010**, *114*, 14450.

(30) Hambourger, M.; Moore, G. F.; Kramer, D. M.; Gust, D.; Moore, A. L.; Moore, T. A. Biology and technology for photochemical fuel production. *Chem. Soc. Rev.* **2009**, *38*, 25.

(31) Bonetti, C.; Stierl, M.; Mathes, T.; van Stokkum, I. H. M.; Mullen, K. M.; Cohen-Stuart, T. A.; van Grondelle, R.; Hegemann, P.; Kennis, J. T. M. The Role of Key Amino Acids in the Photoactivation Pathway of theSynechocystisSlr1694 BLUF Domain. *Biochemistry* **2009**, *48*, 11458.

(32) Berera, R.; van Grondelle, R.; Kennis, J. T. M. Ultrafast transient absorption spectroscopy: principles and application to photosynthetic systems. *Photosynth. Res.* **2009**, *101*, 105.

(33) van Stokkum, I. H. M.; Larsen, D. S.; van Grondelle, R. Global and target analysis of time-resolved spectra. *Biochim. Biophys. Acta, Bioenerg.* 2004, 1657, 82.

(34) Chen, X.; Larsen, D. S.; Bradforth, S. E.; van Stokkum, I. H. M. Broadband Spectral Probing Revealing Ultrafast Photochemical Branching after Ultraviolet Excitation of the Aqueous Phenolate Anion. *J. Phys. Chem. A* **2011**, 230.

## The Journal of Physical Chemistry Letters

(35) Larsen, D. S.; van Stokkum, I. H.; Vengris, M.; van Der Horst, M. A.; de Weerd, F. L.; Hellingwerf, K. J.; van Grondelle, R. Incoherent manipulation of the photoactive yellow protein photocycle with dispersed pump-dump-probe spectroscopy. *Biophys. J.* **2004**, *87*, 1858.

(36) Gauden, M.; Yeremenko, S.; Laan, W.; van Stokkum, I. H. M.; Ihalainen, J. A.; van Grondelle, R.; Hellingwerf, K. J.; Kennis, J. T. M. Photocycle of the flavin-binding photoreceptor AppA, a bacterial transcriptional antirepressor of photosynthesis genes. *Biochemistry* **2005**, *44*, 3653.

(37) Shibata, Y.; Murai, Y.; Satoh, Y.; Fukushima, Y.; Okajima, K.; Ikeuchi, M.; Itoh, S. Acceleration of electron-transfer-induced fluorescence quenching upon conversion to the signaling state in the blue-light receptor, TePixD, from Thermosynechococcus elongatus. *J. Phys. Chem. B* **2009**, *113*, 8192.

(38) Toh, K. C.; van Stokkum, I. H. M.; Hendriks, J.; Alexandre, M. T. A.; Arents, J. C.; Perez, M. A.; van Grondelle, R.; Hellingwerf, K. J.; Kennis, J. T. M. On the signaling mechanism and the absence of photoreversibility in the AppA BLUF domain. *Biophys. J.* **2008**, *95*, 312.

(39) Kennis, J. T. M.; Groot, M.-L. Ultrafast spectroscopy of biological photoreceptors. *Curr. Opin. Struct. Biol.* 2007, 17, 623.

(40) Reece, S. Y.; Hodgkiss, J. M.; Stubbe, J.; Nocera, D. G. Protoncoupled electron transfer: the mechanistic underpinning for radical transport and catalysis in biology. *Philos. Trans. R. Soc. B* **2006**, *361*, 1351.

(41) Song, P. S. On the basicity of the excited state of flavins. *Photochem. Photobiol.* **1968**, *7*, 311.

(42) Fang, C.; Frontiera, R. R.; Tran, R.; Mathies, R. A. Mapping GFP structure evolution during proton transfer with femtosecond Raman spectroscopy. *Nature* **2009**, *462*, 200.

(43) Stelling, A. L.; Ronayne, K. L.; Nappa, J. r. m.; Tonge, P. J.; Meech, S. R. Ultrafast structural dynamics in BLUF domains: transient infrared spectroscopy of AppA and its mutants. *J. Am. Chem. Soc.* **2007**, *129*, 15556.

(44) Unno, M.; Masuda, S.; Ono, T.-a.; Yamauchi, S. Orientation of a key glutamine residue in the BLUF domain from AppA revealed by mutagenesis, spectroscopy, and quantum chemical calculations. *J. Am. Chem. Soc.* **2006**, *128*, 5638.

(45) Grinstead, J. S.; Avila-Perez, M.; Hellingwerf, K. J.; Boelens, R.; Kaptein, R. Light-induced flipping of a conserved glutamine sidechain and its orientation in the AppA BLUF domain. *J. Am. Chem. Soc.* **2006**, *128*, 15066.

(46) Kandori, H.; Iwata, T.; Watanabe, A.; Iseki, M.; Watanabe, M. Strong Donation of the Hydrogen Bond of Tyrosine during Photoactivation of the BLUF Domain. *J. Phys. Chem. Lett.* **2011**, *2*, 1015.

(47) Sadeghian, K.; Bocola, M.; Schütz, M. A QM/MM study on the fast photocycle of blue light using flavin photoreceptors in their light-adapted/active form. *Phys. Chem. Chem. Phys.* **2010**, *12*, 8840–8846.

(48) Götze, J.; Saalfrank, P. Serine in BLUF domains displays spectral importance in computational models. *J. Photochem. Photobiol. B-Biol.* **2009**, *94*, 87.

(49) Yamaguchi, S.; Kamikubo, H.; Kurihara, K.; Kuroki, R.; Niimura, N.; Shimizu, N.; Yamazaki, Y.; Kataoka, M. Low-barrier hydrogen bond in photoactive yellow protein. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 440.