

First Constraint on Cosmological Variation of the Proton-to-Electron Mass Ratio from Two Independent Telescopes

F. van Weerdenburg,^{1,2} M. T. Murphy,³ A. L. Malec,³ L. Kaper,^{1,2} and W. Ubachs¹

¹*Institute for Lasers, Life and Biophotonics, VU University Amsterdam, de Boelelaan 1081, 1081 HV Amsterdam, The Netherlands*

²*Astronomical Institute Anton Pannekoek, Universiteit van Amsterdam, 1098 SJ Amsterdam, The Netherlands*

³*Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Melbourne, Victoria 3122, Australia*

(Received 3 March 2011; published 6 May 2011)

A high signal-to-noise spectrum covering the largest number of hydrogen lines (90 H₂ lines and 6 HD lines) in a high-redshift object was analyzed from an observation along the sight line to the bright quasar source J2123-005 with the Ultraviolet and Visual Echelle Spectrograph on the European Southern Observatory Very Large Telescope (Paranal, Chile). This delivers a constraint on a possible variation of the proton-to-electron mass ratio of $\Delta\mu/\mu = (8.5 \pm 3.6_{\text{stat}} \pm 2.2_{\text{sys}}) \times 10^{-6}$ at redshift $z_{\text{abs}} = 2.059$, which agrees well with a recently published result on the same system observed at the Keck telescope yielding $\Delta\mu/\mu = (5.6 \pm 5.5_{\text{stat}} \pm 2.9_{\text{sys}}) \times 10^{-6}$. Both analyses used the same robust absorption line fitting procedures with detailed consideration of systematic errors.

DOI: 10.1103/PhysRevLett.106.180802

PACS numbers: 06.20.Jr, 33.20.-t, 95.85.Mt, 98.80.Es

Experimental searches for temporal variation of fundamental constants on a cosmological time scale have resurfaced on the agenda of contemporary physics since the ground-breaking study of a possible variation of the fine-structure constant α from quasar observations, now a decade ago [1]. While $\alpha = e^2/4\pi\epsilon_0\hbar c$ is a fundamental constant representing the strength of the electromagnetic force, the proton-to-electron mass ratio $\mu = m_p/m_e$ is another important dimensionless constant to be tested for possible temporal drifts. Since the quark masses contribute only marginally to the proton mass ($\sim 10\%$), m_p is nearly proportional to the strength of the nuclear force Λ_{QCD} . Hence, μ scales with Λ_{QCD}/v_H , v_H being the Higgs vacuum expectation value. Therefore, $\Delta\mu/\mu$ probes the cosmological evolution of the nuclear vs the electroweak sector in the standard model. Interestingly, various models of grand unification predict relations of the form

$$\Delta\mu/\mu = R_{\mu\alpha}\Delta\alpha/\alpha, \quad (1)$$

where the proportionality constant is normally large, i.e., $|R_{\mu\alpha}| \approx 10\text{--}40$ [2], although model-dependent [3]. Measuring any possible changes in α and μ therefore allows a test of “beyond standard” theories.

At cosmological redshifts $z > 2$, molecular hydrogen is the test system for probing a drift in μ [4–6], where the time arrow is defined by $\Delta\mu = \mu_z - \mu_0$. Wavelengths λ^z of spectral lines in the Lyman and Werner bands of H₂ observed in distant objects may be compared to the wavelengths λ^0 for the same lines measured in the laboratory at high accuracy [6–8] via

$$\frac{\lambda_i^z}{\lambda_i^0} = (1 + z_{\text{abs}}) \left(1 + K_i \frac{\Delta\mu}{\mu} \right), \quad (2)$$

with z_{abs} the redshift of the absorbing gas cloud in the line of sight of the quasar and K_i the so-called sensitivity

coefficient, expressing the sensitivity for each spectral line to a mass variation $\Delta\mu$; the K_i can be calculated to an accuracy of 1% [4,6,7]. For the deuterated isotopologue HD, accurate laboratory data and K coefficients are available [9].

Here we report on an investigation of the H₂ absorption system at $z_{\text{abs}} = 2.06$ toward the quasar J2123-005 making use of the Ultraviolet and Visual Echelle Spectrograph (UVES) at the Very Large Telescope (VLT) (European Southern Observatory, Paranal, Chile). The same object was recently investigated with the High Resolution Echelle Spectrometer (HIRES) at the Keck Telescope [10], so that now, for the first time, an accurate μ -variation analysis becomes available from two independent telescopes. Moreover, J2123-005 is the brightest quasar with H₂ absorption, with $R = 15.8$ mag. An 11 h observation in August and September 2008 at VLT delivered a spectrum reaching a signal-to-noise ratio of 65 per 1.5 km s^{-1} pixel at 3415 Å, degrading to ~ 15 at the far UV end (3046 Å). In this range many hydrogen lines are found, of which 90 H₂ and 6 HD are useful to include in an analysis; two typical spectral regions are shown in Fig. 1, while the entire spectrum is reproduced in the supplemental material [11]. Individual spectra (14 exposures of 2900 s) were air-vacuum and heliocentrically corrected, rebinned, and coadded.

This high-quality VLT spectrum of J2123 was subjected to a comprehensive fitting procedure with a direct χ^2 minimization of $\Delta\mu/\mu$ to the entire spectrum, in a similar manner as in Refs. [10,12]. This procedure has many advantages over a line-by-line fitting procedure as used in other studies [7,13,14]. The former allows more objective selection of spectral regions to be fitted, and lines that are partly blended by Lyman- α forest lines can be modeled and explicitly included in the fitting routine. By this means,




FIG. 1 (color online). Spectrum of J2123-005 observed with UVES-VLT clearly showing several H_2 transitions, each comprising two distinct velocity features. Left: Region 3097–3106 Å with a relatively low signal-to-noise ratio. Right: Region 3251–3263 Å toward longer wavelength at a higher signal-to-noise ratio, also containing a line of HD. The upper trace shows residuals in the fit regions. The tick marks above the spectrum refer to the velocity components of the H_2 and HD lines (green), modeled Lyman- α forest lines (blue), and Fe II lines (red). H_2 spectral line identifications are shown at the bottom.

the amount of information on H_2 absorption extracted from the spectrum is effectively increased; however, the regions where no H_2 absorption occurs are cut out of the fitting routine to avoid contaminating the χ^2 . In total, 68 regions cover the 96 H_2 /HD lines included in the fit. Additional regions containing some 39 H_2 /HD lines were excluded, because lines were too weak, were located within a saturated (Lyman- α) absorption region, or were contaminated with unidentified narrow spectral features, i.e., metal lines.

A generic model spectrum is produced from the accurately known transition wavelengths for H_2 [8] and HD lines [9]. Intensities derive from multiplying calculated oscillator strengths [15] with column densities $N(J)$ for each rotational state as fitting parameters. This mask spectrum is then convolved threefold, with a Lorentzian for the damping coefficient, Γ_i , for each line [16], a fixed width parameter for the instrumental profile of the UVES spectrograph (0.8'' resulting in a resolving power of 53 000), and a Gaussian fit parameter b for the Doppler width. In this way, the H_2 and HD lines in the spectrum are tied, leading to a reduction in the number of parameters to be adjusted in the fitting procedure.

This spectrum is replicated to accommodate the velocity structure of the absorbing cloud, starting with the two features split by ~ 25 km/s visible in Fig. 1. Subsequently, additional velocity components (VCs) are added, thereby increasing the number of fit parameters by b_{vc} and $N_{vc}(J)$ for all VCs. Only after a full H_2 /HD velocity structure is imposed, all Lyman-forest and metal lines are assigned and simulated properly, and the continuum and zero levels are adjusted for all local fitting regions is a possible variation of the proton-to-electron mass ratio $\Delta\mu/\mu$ included as a final fitting parameter. The VPFIT algorithm [17] uses Eq. (2) and the K_i coefficients [6,7,10] directly. In addition to Lyman- α forest lines, metal lines falling in the H_2 regions are also cofitted. As one example, we mention the 18 velocity components of the Fe II line, weakly overlapping the H_2 L3R0 line at 3252 Å [11]; its velocity structure is tied in the fit to the Fe II line observed at 7952 Å ($\lambda_{lab} = 2600$ Å).

One of the advantages of fitting all lines simultaneously is that the velocity structure of the absorbing system can be

robustly investigated. For example, a composite residual spectrum was constructed from the 25 cleanest absorption lines, all shifted to a common redshift ($z = 2.059400$), as shown in Fig. 2. It demonstrates that significant H_2 absorption is left in a systematic way in the two-component fit; four- and five-component fits clearly yield better representations of the spectrum. Besides visual inspection of residuals, the procedure of adding velocity components can be quantified by calculating the value of χ^2 per degree of freedom, χ^2_ν , as is shown in Fig. 3. χ^2_ν decreases until a possible 6th VC, which is then rejected on the basis of statistical grounds. Application of a modified version of the Akaike Information Criterion for finite sample sizes yields a value of $\Delta_{AIC} \sim 90$ (for definition and evaluation, see [18]), thus establishing an objective ground for adding a 5th VC and for leaving out the 6th VC. The result of the 5-component fit is thus obtained as the fiducial statistical outcome with $\Delta\mu/\mu = (8.5 \pm 3.6) \times 10^{-6}$.

The robustness of the statistical analysis was further investigated by verifying that the result was not produced by a limited subclass of lines. A fit was performed whereby




FIG. 2 (color online). Composite residual spectrum composed of the 25 cleanest H_2 lines for fits to 2-, 3-, and 5-velocity components. Upper panel: Residuals. Lower panel: The spectrum as simulated from the final parameters resulting in the 2- and 5-VC models. The tick marks indicate the fitted positions in 2- and 5-VC models; in the 5-VC model, two VCs lie at almost the same velocity and appear as a single tick mark on this plot.

a different value of $\Delta\mu/\mu$ was fitted simultaneously to each of the rotational states in hydrogen, $H_2(J)$, and for HD. The results shown in Fig. 3(c) demonstrate the consistency for the value of $\Delta\mu/\mu$ in this procedure. In the model description, the physical assumption was made that the Doppler widths b_{vc} (for each velocity component z_{vc}) were independent of the rotational state; if, however, a fit is performed with varying Doppler widths $b_v(J)$, a $\Delta\mu/\mu$ value is returned differing by only 5%.

The analysis strategy aimed to stick as close as possible to the known molecular physics of H_2 and HD. The damping parameters Γ_i from Ref. [16] describe the wings of the saturated lines (as in Fig. 1) well. The oscillator strengths f_i were kept fixed to the molecular physics values [15], thereby keeping the number of fit parameters as low as possible. However, in the region 3345–3415 Å, covering the L0 and L1 Lyman lines, the calculated f_i underestimate the line intensities by some 50%. Hence, in the fits the oscillator strengths in this range were adjusted, similarly as in Ref. [10], where this phenomenon was attributed to the O IV/Lyman- β emission feature of the background quasar.

As for systematic effects on the resulting value of $\Delta\mu/\mu$, adverse influences on the wavelength calibration are of particular importance. Wavelength calibration of the composite spectrum was derived from Th-Ar frames attached to each science exposure, where use was made of the improved Th-Ar line selection algorithm rejecting blended and weak lines [19]. Wavelength calibration residuals of Th-Ar lines, used for calibrating the echelle grating, are typically 70 m/s. However, since the wavelength scale of each echelle order is derived from typically more than 11 Th-Ar lines and many H_2 transitions are fitted for each order, this effect largely averages out, and long-range variations of the wavelength scale are limited to 30 m/s [19]. In view of a span in K_i of ~ 0.05 , a shift of 30 m/s translates into a maximum variation in $\Delta\mu/\mu$ of

2.0×10^{-6} ; here, velocity shifts are expressed as $\Delta v = cK_i\Delta\mu/\mu$.

In addition, intraorder distortions within individual echelle orders, first studied for the HIRES-Keck grating system [20] and later for UVES-VLT [21], may result in peak-to-peak velocity shifts of up to $\Delta v = 200$ m/s between transitions. This phenomenon was simulated by producing a counterdistorted spectrum by shifting the center of all echelle orders by -100 m/s and imposing shifts up to $+100$ m/s at the order edges. From a fit to this spectrum, it is estimated that this phenomenon produces a possible systematic effect of 0.7×10^{-6} . Systematic effects from spectral redispersion occurring in the rebinning process when averaging over the 14 science exposures are less than 0.2×10^{-6} . Finally, drifts in temperature and atmospheric pressure between the quasar and Th-Ar calibration exposures are <1 K and <1 mbar, respectively, leading to a possible error of $<0.7 \times 10^{-6}$. By taking the possible shifts in quadrature, this results in an estimate of the systematic error of $\delta_{\text{sys}}(\Delta\mu/\mu) = 2.2 \times 10^{-6}$. Uncertainties in the K_i coefficients may lead to uncertainties in $\Delta\mu/\mu$ at the $\sim 0.01 \times \Delta\mu/\mu$ level [6]; i.e., they are negligible.

A $\Delta\mu$ effect can be mimicked by a long-range distortion of the wavelength scale due to the correlation of K coefficients with wavelength [6]. Separate fitting of the range $\lambda < 3230$ Å, where Werner and Lyman lines exhibiting strongly differing K_i values are found alongside each other, returns a value within 10% of the fiducial value, although with a larger uncertainty margin. This can be considered as an internal consistency check of the wavelength scale.

The statistical and systematic analyses lead to a final result for the UVES-VLT H_2 absorption spectrum toward J2123 (uncertainties representing 1σ):

$$\Delta\mu/\mu = (8.5 \pm 3.6_{\text{stat}} \pm 2.2_{\text{sys}}) \times 10^{-6}. \quad (3)$$

This may be compared to the result from the analysis of the HIRES-Keck spectrum for the same object [10]:

$$\Delta\mu/\mu = (5.6 \pm 5.5_{\text{stat}} \pm 2.9_{\text{sys}}) \times 10^{-6}. \quad (4)$$

These values are well within 1σ of each other and may be averaged to yield $\Delta\mu/\mu = (7.6 \pm 3.5_{\text{tot}}) \times 10^{-6}$. The agreement obtained between both investigations of the same high-redshift H_2 absorption spectrum from the two major large telescopes VLT and Keck enhances the confidence in the statistical and systematic uncertainty analyses. They are indicative of the error sources being suitably taken into account for both instruments.

Spectral observations of highly redshifted H_2 absorbers are limited. Of the many thousands of known quasar systems, of which some 1000 have been classified as damped-Lyman- α systems, in only some 20 is molecular hydrogen detected but not under conditions allowing for a detailed analysis of μ variation at a competitive level. So far, only




FIG. 3 (color online). Results from various fits. (a) Resulting value for $\Delta\mu/\mu$ for progressively increasing the number of velocity components; there are two results for 3-VC fits, obtained by putting the 3rd component in both distinct velocity features, which are horizontally offset for clarity; (b) the χ^2_{ν} value; (c) result on $\Delta\mu/\mu$ from fits to 7 subclasses of spectral lines.




FIG. 4 (color online). Current constraints on $\Delta\mu/\mu$ on a cosmological time scale. Black points for $z < 1$ refer to NH_3 values from objects B0218 [25,26] and PKS1830 [27]; blue diamonds refer to Ref. [12]; and the green triangle to Ref. [10], i.e., the HIRES-Keck analysis of J2123. The gray circle refers to a recent reanalysis of Q0347 [22]. The red square is the present result of our UVES-VLT analysis of J2123. All uncertainties are at 1σ . Independent constraints on the same object have been offset in redshift for clarity.

high-quality spectra from H_2 absorbing systems toward Q0347-383 at $z_{\text{abs}} = 3.02$ and Q0405-443 at $z_{\text{abs}} = 2.59$ [7,12] and Q0528-250 at $z_{\text{abs}} = 2.81$ [12] have been observed with the Very Large Telescope to yield a tight constraint on $\Delta\mu/\mu$. Recently, a VLT reinvestigation of Q0347-383 was also reported [22]. In contrast, for detecting drifts in α , spectra of atomic species (Mg I, Mg II, Fe II, Zn II, Cr II, Si II, etc.) in almost 300 absorption systems have been analyzed [23]. For H_2 , a spectrum of J1337 + 315 at $z_{\text{abs}} = 3.17$ was analyzed [24], but due to the low column density [$\log N(\text{H}_2) = 14.1$] a constraint on $\Delta\mu/\mu$ only at the noncompetitive 10^{-4} level was found.

In Fig. 4, the now existing information on high-redshift absorbing H_2 systems is collected; the tight constraints from the ammonia method [25–27] for $z < 1$ are included as well. The status result of a possible variation of the proton-to-electron mass ratio can be expressed as follows: There exists no firm evidence for a drifting μ on a cosmological time scale, which is constrained by $\Delta\mu/\mu < 1 \times 10^{-5}$ at redshifts in the range $z = 2$ –3. Quantitatively, averaging the six results for the four high-redshift H_2 absorption systems yields $\Delta\mu/\mu = (5.2 \pm 2.2) \times 10^{-6}$, which is a slight indication (at 2.3σ level) for a larger μ at high redshift, but further systems should be analyzed to warrant such a claim. As for the proportionality relation of Eq. (1), combined with the finding of $\Delta\alpha/\alpha = (-5.4 \pm 1.2) \times 10^{-6}$ [23], some significant drift on μ exceeding the present constraint would be expected. The result may also be interpreted as the specific assumptions of grand unification underlying Eq. (1) to be unjustified. Indeed, while the majority of the grand unified theory scenarios predict $|R_{\mu\alpha}|$ to be large, some special cases

are reported yielding a smaller proportionality constant, i.e., $|R_{\mu\alpha}| \leq 5$ [3]. The present results constraining $\Delta\mu/\mu$ may thus be used to discriminate between various possible grand unified theory scenarios.

This work is based on observations carried out at the European Southern Observatory under program No. 81.A-0242 (PI Ubachs), with the UVES spectrograph installed at the Kueyen UT2 on Cerro Paranal, Chile. The authors acknowledge fruitful discussions with J. King and J. K. Webb (Sydney). M. T. M. thanks the Australian Research Council, and W. U. thanks the Netherlands Foundation for Fundamental Research on Matter (FOM) for research support.

-
- [1] J. K. Webb *et al.*, *Phys. Rev. Lett.* **82**, 884 (1999).
 - [2] V. V. Flambaum *et al.*, *Phys. Rev. D* **69**, 115006 (2004); X. Calmet and H. Fritsch, *Eur. Phys. J. C* **24**, 639 (2002).
 - [3] T. Dent, S. Stern, and C. Wetterich, *Phys. Rev. D* **78**, 103518 (2008).
 - [4] R. Thompson, *Astrophys. Lett. Commun.* **16**, 3 (1975).
 - [5] D. A. Varshalovich and S. A. Levshakov, *JETP Lett.* **58**, 237 (1993).
 - [6] W. Ubachs *et al.*, *J. Mol. Spectrosc.* **241**, 155 (2007).
 - [7] E. Reinhold *et al.*, *Phys. Rev. Lett.* **96**, 151101 (2006).
 - [8] E. J. Salumbides *et al.*, *Phys. Rev. Lett.* **101**, 223001 (2008); D. Bailly *et al.*, *Mol. Phys.* **108**, 827 (2010).
 - [9] T. I. Ivanov *et al.*, *Phys. Rev. Lett.* **100**, 093007 (2008).
 - [10] A. L. Malec *et al.*, *Mon. Not. R. Astron. Soc.* **403**, 1541 (2010).
 - [11] See supplemental material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.106.180802> for the full spectrum of VLT-J2123.
 - [12] J. A. King *et al.*, *Phys. Rev. Lett.* **101**, 251304 (2008).
 - [13] A. Ivanchik *et al.*, *Astron. Astrophys.* **440**, 45 (2005).
 - [14] R. I. Thompson *et al.*, *Astrophys. J.* **703**, 1648 (2009).
 - [15] H. Abgrall *et al.*, *Can. J. Phys.* **72**, 856 (1994).
 - [16] H. Abgrall, E. Roueff, and I. Drira, *Astron. Astrophys. Suppl. Ser.* **141**, 297 (2000).
 - [17] <http://www.ast.cam.ac.uk/~rfc/vpfit.html>.
 - [18] N. Sigiura, *Commun. Stat., Theory Methods* **7**, 13 (1978).
 - [19] M. T. Murphy *et al.*, *Mon. Not. R. Astron. Soc.* **378**, 221 (2007).
 - [20] K. Griest *et al.*, *Astrophys. J.* **708**, 158 (2010).
 - [21] J. B. Whitmore, M. T. Murphy, and K. Griest, *Astrophys. J.* **723**, 89 (2010).
 - [22] M. Wendt and P. Molaro, *Astron. Astrophys.* **526**, A96 (2011).
 - [23] M. T. Murphy, J. K. Webb, and V. V. Flambaum, *Mon. Not. R. Astron. Soc.* **345**, 609 (2003).
 - [24] R. Srianand *et al.*, *Mon. Not. R. Astron. Soc.* **405**, 1888 (2010).
 - [25] M. T. Murphy *et al.*, *Science* **320**, 1611 (2008).
 - [26] N. Kanekar, *Astrophys. J. Lett.* **728**, L12 (2011).
 - [27] C. Henkel *et al.*, *Astron. Astrophys.* **500**, 725 (2009).